

Dr. Vijay

①

ISSN 2277 - 5730
AN INTERNATIONAL MULTIDISCIPLINARY
QUARTERLY RESEARCH JOURNAL

AJANTA

Volume - XII Issue - I

January - March - 2023

ENGLISH PART II / MARATHI

Peer Reviewed Refereed
and UGC Listed Journal
Journal No. 40776



ज्ञान-विज्ञान विमुक्तये

IMPACT FACTOR / INDEXING
2020 - 6.306
www.sjifactor.com

❖ EDITOR ❖

Asst. Prof. Vinay Shankarrao Hatole

M.Sc (Maths), M.B.A. (Mktg.), M.B.A. (H.R.),
M.Drama (Acting), M.Drama (Prod. & Dir.), M.Ed.

❖ PUBLISHED BY ❖

Ajanta Prakashan
Aurangabad. (M.S.)

Handwritten signature

VC Principal

Reena Mehta College of Arts, Science

Commerce & Mgt. Studies

Bhayandar (West), Dist. Thane - 401 10



THE UNIVERSITY OF CHICAGO
LIBRARY

❧ CONTENTS OF ENGLISH PART - II ❧

S.No.	Title & Author	Page No.
1	Perception of undergraduates towards transit advertising Ms. Subhashini Iyengar Dr. Pooja Ramchandani	1-12
2 ✓	Paraben Derivatives as Future Potential Drug Vijay D. Gangan Uttam Yadav Kundan Mewad Yogita Purohit Neha Rastogi Priyanka A. Save Anamika Singh Krishnalekha Nadar	13-21 13-22
3 ✓	To Study Investment Patterns and Customer Attitudes in Relation to Mutual Funds Ms. Abeda Shaikh Dr. Vaishali Kothiya	22-32 23-33
4	The Dynamics of Sustainable Development: Social change through Teacher Education Programme Dr. Shaheen Parveen	33-38
5	Impact of Technology on Different Leadership Styles to Support Distanced Working: A Study of India's Service Sector in Telangana State Linga Reddy Katipalli	39-45
6	Challenges of Human Resource Management in Current Scenario Ms. Saniya Shiurkar U. Dr. Padmapani Bhagwan Sawai	46-50
7 ✓	Production of Cost Effective, Biodegradable, Disposable Sanitary Napkins using Banana Fibres Dr. Vijay D. Gangan Dr. Uttam Yadav Yogita Purohit Neha Rastogi Krushnal Devlekar Mrs. Priyanka A. Save	51-58 52-58




IC Principal
 Reena Mehta College of Arts, Science
 Commerce & Mgt. Studies
 Bhayandar (West), Dist. Thane - 401 101

❧ CONTENTS OF ENGLISH PART - II ❧

S. No.	Title & Author	Page No.
8	Traditional Accounting and Modern Accounting system in India with Special Reference to Frauds after 2000-01 Shaikh Mumtaz	59-667
9	Digital Transformation in the Economy is a Boon! Dr. Prakash Laxmanrao Dompale	68-74
10 ✓	A Study on Students' Attitudes towards Online Shopping Platforms Amit Chhotelal Gupta Urmila Arjun Chauhan	75-81
11 ✓	Statistics and Blood Glucose Monitoring Krishnalekha Nadar Vijay D. Gangan Priyanka A. Save Sukanya Sawant	21-86 82-86
12 ✓	Subliminal Advertising : Utilizing Deception to Influence Consumption Habits Mr. Frank Remedios	87-92
13	Higher Initiative to Learn: Essential Quality for Entrepreneurship Development Prof. Himanshul Panchal	93-96
14	Game Changer: The Role of Sports in Achieving Sustainable National Development Dr. Surayakant Pandurand Dhonde	97-101

2. Paraben Derivatives as Future Potential Drug

Vijay D. Gangan

Department of Chemistry, Reena Mehta College, Bhayandar (W), Mumbai.

Uttam Yadav

Kundan Mewad

Department of Chemistry, Andheri Bhawan's College, Mumbai.

Yogita Purohit

Neha Rastogi

Priyanka A. Save

Anamika Singh

Krishnalekha Nadar

Department of Chemistry, Reena Mehta College, Bhayandar (W), Mumbai.

Abstract

Parabens are class of chemicals widely used as preservatives in the cosmetic and pharmaceutical industries. They are attractive preservatives in many types of formulas. These compounds and their salts are used primarily for their bacterial and fungicidal properties. They are also used as food additives. Their analogues also possess various biological activities which prompted us to synthesize few more analogues for their future application as bioactive molecules. All the synthesized compounds were characterized by ¹HNMR and mass spectral data and screened for their antibacterial activity against Gram + ve and Gram – ve cultures. Few of them are showing promising antibacterial activity.

Keywords : Parabens, antibacterial, fungicidal, ¹HNR, elemental analysis.

Introduction

Phenolic phytochemicals are known to exhibit anti-inflammatory, antioxidant, anticarcinogenic, antidiabetic, antiatherosclerosis and immunomodulatory activities in animals¹, 2. These are mostly polyphenols known as secondary plant metabolites³ present in plant and trees. One of such compound is 4-hydroxy benzoic acid which is used as antifungal, antimutagenic, antisickling, esterogenic⁴ and antimicrobial⁵ agent. It is primarily known as the basis for the preparation of its esters, known as parabens, which are used as preservatives in cosmetics. Parabens are used for their bactericidal and fungicidal properties. They can be found in shampoos, commercial moisturizers, shaving gels, personal lubricants, topical / parenteral

pharmaceuticals, spray tanning solution, makeup and toothpaste. They are also used as food additives. In the present study, we are converting 4-hydroxy benzoic acid to methyl and ethyl paraben using conventional method and their further diversification to ester derivatives. Since methyl and ethyl paraben are naturally occurring active compounds having antioxidant and antimicrobial properties, we decided to make a library of compounds using various permutation and combinations to come up with novel ether and ester derivatives of methyl and ethyl paraben using conventional methods. The objective of this study is to condense two molecules of the same disease domain to produce more potent candidate in the same disease domain or to condense two molecules of different disease domain to produce mixed variety of those disease domain or to have drug candidate with entirely different disease domain. In the present work, we are converting 4-hydroxy benzoic acid to methyl and ethyl paraben which in turn further converted to ether and hybrid derivatives respectively using conventional methods.

Results and Discussion

Preparation of methyl and ethyl paraben

They were prepared by refluxing 4-hydroxy benzoic acid with methanol / ethanol using sulphuric acid as a catalyst for 8 hrs. The progress of the reaction is monitored by TLC for the completion of reaction.

Work up

The reaction mixture concentrated under reduced pressure to minimum and to that 200 ml of dichloromethane + 200 ml of water was added. The aqueous layer was extracted successively with dichloromethane (2 x 100 ml). The total organic layer was washed with water (200 ml), brine (100 ml) and concentrated to yield methyl and paraben respectively which can directly used for further diversification. The general yields were 95 – 98 %

General method for the preparation of compounds (I and II)


These were prepared by following general method as depicted below.

To a stirred solution of [A] (1 eq.) in 30 ml acetone was added [B] (2.5 eq.) and stirring continued at 400 C for the next 30 min. for complete formation of K-salt. To this compound [C] (2 eq.) was added and stirring continued at 45-50° C for the next 8 hrs. The progress of the reaction is monitored by TLC for the completion of this reaction.

Work Up

The reaction mixture filtered through Buchner funnel, wash the cake with 25 ml acetone. The total organic layer was concentrated to minimum, preadsorbed on silica gel and purified by silica gel (100 - 200 mesh) column chromatography with increase in concentration of ethyl acetate in petroleum ether. The general yields ranges between 60-70 %.

Reaction Scheme :




Ethyl 4-hydroxybenzoate

Ethyl paraben ether derivatives

Compound No.	R
I	Heptyl
II	3-Methyl butyl

Taking Isoeugenol as general example, the probable mechanism for ethers can be given as follows.



Compound I Ethyl 4 - heptoxybenzoate.

¹H NMR (CDCl₃, 400 MHz) δ ppm : 0.88 (t, J = 4.7 Hz, 3H, terminal -CH₃ from heptoxy moiety), 1.2 - 1.5 (m, 8H, 4 x -CH₂, methylenes from n- heptyl moiety), 1.37 (t, J = 6.9 Hz, 3H, from -OCH₂CH₃ group) , 1.7 - 2.0 (m, 2H, 1x -CH₂ from n-heptyl moiety), 3.99 (t, J = 6.6 Hz, 2H, 1 x -OCH₂ from n-heptyl moiety), 4.34 (q, J = 7.7 Hz, 2H, -OCH₂ from -OCH₂CH₃ group), 6.89 (d, J = 8.7 Hz, 2H, ArH), 7.97 (d, J = 8.7 Hz, 2H, ArH) Molecular Formula C₁₆H₂₄O₃ Pure viscous mass (0.954 gms, 60 %). Anal. Calcd. for C₁₆H₂₄O₃ : C 72.69, H 9.15, O 18.16 Found C 72.66 , H 9.12, O18.19

Compound II Ethyl 4- isopentyloxybenzoate

¹H NMR(CDCl₃, 400 MHz) δ ppm- 0.98 (d, J = 6.6 Hz, 2 x -CH₃, 6H), 1.37 (t, J = 6.9Hz, 3H from -OCH₂CH₃ group), 1.7 (m, 2H, -CH₂ from 3-Methyl butane moiety), 1.85 (s, 1H, CH(CH₃)₂), 4.0 (t, J = 6.6Hz, 2H, 1 x -OCH₂ from 3-Methyl butane moiety), 4.34 (q, J = 7.7 Hz, 2H, -OCH₂ from ethyl paraben moiety), 6.89 (d, J = 8.7 Hz, 2H, ArH), 7.07 (d, J = 8.7 Hz, 2H, ArH). Molecular Formula C₁₄H₂₀O₃. Pure viscous mass (0.894 gms, 63 %). Anal. Calcd .for C₁₄H₂₀O₃ : C 71.16, H 8.53, O 20.31 Found C 71.13, H 8.50, O20.34


General method for the preparation of **compounds (III - V)** These were prepared by following general method as depicted below.

To a stirred solution of Ethyl paraben (1 eq.) in dichloromethane (30 ml) was added pyridine (2.5 eq.) and cool the reaction in ice bath at 15° C. Clear solution of reaction mixture was obtained. To this, was added Acetyl chloride/ benzoyl chloride / substituted benzoyl chloride (2 eq.) at 15 - 20° C and stirred, allowed to attain the room temperature and stirring was continued for the next 24 hrs. (TLC). The organic layer was concentrated under reduced pressure to minimum, preadsorbed on the silica gel and purified by column chromatography (SiO₂, 100-200 mesh) with increase in concentration of Ethyl acetate in petroleum ether to yield pure compound. The purified compounds were unambiguously characterized by ¹HNMR and mass spectroscopy. The general yields of these reactions were ranges between 60 - 80 %.

The most significant features of this methodology are (a) good accessibility of the reagents and its stability (b) a stoichiometric amount of reagent can be used by direct weighing, avoiding excess (c) no evolution of hazardous vapors during the reaction (d) the total elimination of the use of toxic organic solvents (e) a simple experimental procedure (g) good control over the outcome of the reaction by varying the amount of reagent. The aforesaid protocol thus provides

an improved procedure for the synthesis of useful paraben derivatives having important pharmaceutical, agricultural and other physicochemical properties.

Reaction Scheme :




Ethyl 4-hydroxybenzoate

Ethyl paraben ester derivatives

Compound No.	R
III	Acetyl
IV	Benzoyl
V	4-Chloro benzoyl

The probable mechanism for esters can be given as follows.

Probable mechanism for Esters :



Compound III

Ethyl 4 - acetoxybenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm : 1.3 (t, J = 7.7 Hz, 3H, from -OCH₂CH₃), 2.31 (s, 3H, from >COCH₃ group), 4.37 (q, J = 7.14 Hz, 2H, from -OCH₂CH₃), 7.16 (d, J = 8.4 Hz, 2H, ArH), 8.07 (d, J = 8.4 Hz, ArH). Molecular Formula C₁₁H₁₂O₄. Pure viscous mass (0.90 gms, 72 %). Anal. Calcd. for C₁₁H₁₂O₄ : C 63.45, H 5.81, O 30.74 Found C 63.41, H 5.78, O 30.78

Compound IV

Ethyl 4 - benzoyloxybenzoate


¹H NMR (CDCl₃, 400 MHz) δ ppm : 1.4 (t, J = 4.7 Hz, 3H, -CH₃ from -OCH₂CH₃ group), 4.34 (q, J = 7.7 Hz, 2H, -OCH₂ from Ethyl paraben moiety), 7.2 - 8.4 (m, 9H, Ar-H). Molecular Formula C₁₆H₁₄O₄. Off white solid (0.975 gms, 64 %); Melting range 65 - 680C; Anal. Calcd. for C₁₆H₁₄O₄ : C 71.10, H 5.22, O 23.68 Found C 71.06, H 5.19, O 2

Compound V

Ethyl 4-(4-chlorobenzoyl) oxybenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm : 1.4 (t, J = 4.7 Hz, 3H, -CH₃ from -OCH₂CH₃ group), 4.34 (q, J = 7.7 Hz, 2H, OCH₂ from ethyl paraben moiety), 7.0 - 8.7 (m, 8H, ArH). Molecular Formula C₁₆H₁₃ClO₄. Off white solid (1.34 gms, 74 %). Melting range 100 - 1040C Anal. Calcd. for C₁₆H₁₃ClO₄ : C 63.06, H 4.30, O 21.00 Cl 11.63 Found C 63.10, H 4.26, O 21.03

General method for the preparation of compounds (VI to IX):- These were prepared by the general method as mentioned for ethyl paraben.

Reaction Scheme :

Compound No.	R
VI	Acetyl
VII	Benzoyl
VIII	4-Chloro benzoyl
IX	2-Chloro Benzoyl

The general mechanism for ester is same as depicted for ethyl paraben.

Compound VI

Methyl 4-acetoxybenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm : 2.3 (s, 3H, >CO-CH₃), 3.9 (s, 3H, -COOCH₃), 7.16 (d, J = 8.4 Hz, 2H, ArH), 8.07 (d, J = 8.4 Hz, ArH). Molecular Formula C₁₀H₁₀O₄; White solid (1.297 gms, 68 %). Melting range 70 - 750C. Anal. Calcd .for C₁₀H₁₀O₄: C 61.85, H 5.19, O 32.96 Found C 61.81, H 5.16, O 32.99

Compound VII

Methyl 4-benzoyloxybenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm: 3.9 (s, 3H, from -COOCH₃ group), 7.22 (d, J = 8.4 Hz, 2H, ArH), 8.07(d, J = 8.4 Hz, ArH from methyl paraben moiety), 7.2 - 8.3 (m, 5H, ArH from Benzoyl Moiety). Molecular Formula C₁₅H₁₂O₄; White solid (1.09 gms, 65 %). Melting range 106 - 1100C Anal. Calcd .for C₁₅H₁₂O₄ C 70.31, H 4.72, O 24.97. Found C 70.34, H 4.69, O 24.95.

Compound VIII

Methyl 4-(4-chlorobenzoyl)oxybenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm : 3.9(s, 3H,-OCH₃ from -COOCH₃ group), 7.22(d, J = 8.4 Hz, 2H, ArH), 8.1 (d, J = 8.4 Hz, 2H, ArH from Methyl paraben moiety), 7.2 - 8.2 (m, 4H, ArH from 4-Chloro benzoyl moiety). Molecular Formula C₁₅H₁₁ClO₄. Off white solid (1.297 gms, 68 %). Melting range 126 - 1300C. Anal. Calcd .for C₁₅H₁₁ClO₄ : C 61.98, H 3.81, O 22.02, Cl 12.20 Found C 61.95, H 3.78, O 22.05

Compound IX

(4- methoxycarbonyl phenyl)-2-chlorobenzoate

¹H NMR (CDCl₃, 400 MHz) δ ppm : 3.9 (s, 3H,-OCH₃ from -COOCH₃ group), 7.24 (d, J = 8.4 Hz, 2H, ArH from Methyl paraben moiety), 8.1 (d, J = 8.4 Hz, H, ArH from methyl paraben moiety), 7.2 - 8.2 (m, 4H, ArH from 2-Chloro benzoyl moiety). Molecular Formula

C₁₅H₁₁ClO₄. White solid (1.297 gms, 68 %). Melting range 67 - 700C Anal. Calcd for C₁₅H₁₁ClO₄ C 61.98, H 3.81, O 22.02 Cl 12.20 Found C 61.95 , H 3.78 , O 22.03

Chromatographic system

Column chromatography: For column chromatography 100 – 200 mesh Acme grade silica gel is used. The crude reaction mixture is concentrated under reduced pressure to yield crude mass which is preadsorbed on silica gel and purified by column chromatography with increase in concentration of Ethyl acetate in Petroleum ether. The fractions having similar 'rf' values were pooled together, concentrated and subjected for characterization using various spectroscopic techniques.

Thin layer chromatography TLC plates were prepared using silica gel G (ACME, BOMBAY). Pet. ether: EtOAc (85: 15) was used as the solvent system.

Radial chromatography The circular glass plates of thickness 1 mm, were prepared by using silica gel (PF254, E. MERCK, 50 g) in cold distilled water (105 ml). For elution, gradually increasing concentrations of EtOAc in pet ether were employed.

Biological Activity

Antibacterial activity using agar diffusion method 6 Concentration 100 µm

The synthesized molecules were screened for their antibacterial activity using agar diffusion method at 100 µm concentration against Gram positive (*Staphylococcus aureus*) and Gram negative (*Escherichia coli*) bacterial species qualitatively. The results of the antibacterial activities are summarized in Table 1.

Table 1: Antibacterial Activity Results

Sr. No	Compound No.	Antibacterial Activity	
		Against Gram - ve bacteria species (<i>Escherichia coli</i>)	Against Gram +ve bacterial species (<i>Staphylococcus aureus</i>)
1	Methyl paraben	+	+
2	Ethyl paraben	-	-
3	Ampiciline	+	+

	(Std. Drug)		
4	I	-	-
4	II	+	-
5	III	-	-
6	IV	-	-
7	V	-	+
8	VI	-	-
9	VII	-	-
10	VIII	-	+
11	IX	-	-

The above results shows that the base molecule Methyl paraben has anti-bacterial activity against both the bacterial culture. Its derivative viz. V and VIII were active against *Staphylococcus aureus* (Gram + ve bacteria) and II was active against *Escherichia coli* (Gram - ve bacteria) respectively. Thus, 3-methyl butyl and 4-chloro benzoyl derivatives were potential antibacterial candidates. In depth analysis of these compounds through structure activity relationship studies would provide further insight and can be an interesting topic of future studies.

The structural diversity and the pronounced biological activities encountered in the paraben derivatives suggests that this class of compounds is worthy for further studies that may lead to derivatives by using combinatorial chemistry approach is an alternative strategy to new therapeutic discovery. In other words the generation of diverse benzofuran derivatives develop new therapeutic molecules that might result in candidates having better activity.

References

1. Wattenberg L. W., Coccia J. B., Lam L. K. Inhibitory effects of phenolic compounds on benzo(a)pyrene-induced neoplasia. *Cancer Res.*, 1980, 40, pp. 2820 – 2823.



VC Principal
Reena Mehta College of Arts, Science,
Commerce & Mgt. Studies

2. Talalay P., De Long M. U., Prochaska H. J. Identification of a common chemical signal regulating the induction of enzymes that protects against chemical carcinogenesis. *Proc. Natl. Acad. Sci. USA*.
3. Macheix JJ., Fleuriet A.;Billot J. Fruit phenolics. Boca Raton, FL; CTC; 1990.
4. Pugazhendhi, D., Pope G. S., Darbre, P. D. Oestrogenic activity of p-hydroxy benzoic acid and methyl paraben in human breast cancer cell lines. *J. Appl. Toxicol.* 2005, 25, pp. 301 – 309.
5. Chong K. P., Rossall S.' Atong M. In vitro antimicrobial activity and functioning of syringic acid, caffeic acid and 4-hydroxy benzoic acid against ganoderma boninense. *J. Agr. Sci.* 2009, 1, pp. 15 – 20.
6. Finn R. K. Theory of Agar Diffussion Methods for Bioassay. *Anal. Chem.*, 1959, 31 (6), pp 975.